



Optical tracking of ultrafast surface vibrations

Wright OB^{(1)*}

(1) Hokkaido University, Sapporo 060-8628, Japan

*Corresponding author's email: olly@eng.hokudai.ac.jp

Tracking phonons at surfaces is crucial to understanding the internal structure of solids. This talk describes how to do this by spatiotemporal mapping of surface motion at frequencies from 100 MHz up to above 100 GHz, giving a new eye for subsurface features down to the atomic scale as well as revealing artistic phonon fields in crystals, cavities and waveguides.

References

- [1] O.B. Wright, K. Kawashima, Coherent phonon generation from ultrafast surface vibrations, Phys. Rev. Lett. 69 (1992) 1668.
- [2] Y. Sugawara et al., Watching ripples on crystals, Phys. Rev. Lett. 88 (2002) 185504.
- [3] D. M. Profunser et. al., Imaging ripples on phononic crystals reveals acoustic band structure and Bloch harmonics, Phys. Rev. Lett. 97 (2006) 055502.
- [4] T. Tachizaki et al., Acoustic whispering-gallery modes generated and dynamically imaged with ultrashort optical pulses, Phys. Rev. B 81 (2010) 165434-1-5.
- [5] P. H. Otsuka et al., Broadband evolution of phononic-crystal-waveguide eigenstates in real- and k-spaces, Sci. Rep. 3 (2013) 3351.
- [6] P. H. Otsuka et al., Time-domain imaging of gigahertz surface waves on an acoustic metamaterial, New J. Phys. 20 (2018) 013026.
- [7] Q. Xie et al., Imaging GHz zero-group-velocity Lamb waves, Nat. Comm. 10 (2019) 2228.